

## Initiation and Engagement of Alcohol and Other Drug Use Treatment (IET)

Performance Measure Set:  CCO Incentive  Medicare Star Rating

Quality Measurement Type:  Structure  Process  Outcome  Patient Experience

Data Type:  Claims  Chart Documentation  eCQM  Survey  Other

State Benchmark: Initiation for Age 18+ – 46.8% (2019 National Medicaid 75th percentile)

Engagement for Age 18+ – 18.5% (2019 National Medicaid 75th percentile)

*Must meet both components to achieve measure.*

**Who:** Members aged 18 years and older with a new diagnosis of alcohol or other drug use between January 1, 2021–November 14, 2021. A diagnosis is considered “new” if the member has not had a diagnosis of (or received medication for) alcohol or other drug use in the previous 60 days.

**Why:** Access to treatment for substance use disorder is a critical aspect of a person’s health and their journey through recovery. The IET metric is a tool to encourage coordination across the network of care providers for substance use treatment and helps ensure people have timely access to appropriate care.

**What:** Two rates are reported for this measure: Initiation and Engagement. Both measures use the same denominator.

1. Initiation – For members with a new episode of alcohol or other drug use (diagnosis on a claim with no other diagnosis in the previous 60 days), this metric measures the percentage of those who initiated treatment within 14 days through either medication dispensing or a SUD visit with a provider.
  - a. Initiation of treatment can be on the same day as the new alcohol or other drug use diagnosis if the services are with different providers.
2. Engagement - For members with a new episode of alcohol or other drug use (diagnosis on a claim with no other diagnosis in the previous 60 days), this metric measures the percentage of those who had two treatment events, either medication dispensing or a SUD visit with a provider, within 34 days from their initial treatment event.
  - a. If treatment was initiated through a medication dispensing event, only one of the two required engagement events can be through medication and the other must be through a SUD visit with a provider.
  - b. Both engagement events can be on the same day if the services are with different providers; the exception being if one event is for medication-assisted treatment there is no requirement that they be different providers.

**How:** There are over 230 codes that count toward numerator criteria through a visit with a provider; **please see IET Guide for Primary Care on the following pages for additional details.** In general, initiation and engagement events can be through medication dispensing events, inpatient, outpatient, observation, or telemedicine visits.

NOTE: Methadone is not included in the medication lists for this measure because Methadone for opioid use disorder does not show up in pharmacy claims data. However, Methadone for opioid use disorder treatment **does count as treatment for this metric** and would be captured on medical claims.

**Exclusions:** Hospice during any point in the year.

## Initiation and Engagement of Alcohol and Other Drug Use Treatment (IET) FAQs

**Q: Is tobacco use included in this metric?**

**A:** No. While we do consider tobacco use disorder to be included in the continuum of substance use disorders from a clinical perspective, it is not considered as one of the diagnosis codes that would qualify someone for the IET metric.

**Q: What is considered as “other drugs” in this metric?**

**A:** The IET measure is looking for substance use disorder diagnosis including alcohol, opioid and other drugs such as cocaine, cannabis, methamphetamine, hypnotics, sedatives, inhalants, etc. See OHA specifications for full list.

**Q: How are initial alcohol or other drug use diagnoses identified?**

**A:** Alcohol or other drug use disorder diagnosis codes are identified using claims for services that occurred in the following visit types:

- Outpatient visits
- Telehealth
- E-visit of virtual check-in
- Intensive outpatient visits
- Partial hospitalization
- Detoxification visits
- ED visits or Observation
- Acute or non-acute inpatient admits
- Online assessment
- Opioid treatment services

## IET Guide for Primary Care

*This guide includes 4 sections:*

- 1) Billing and Coding
- 2) BHC engagement
- 3) Collective Medical - IET cohort, useful tips
- 4) Person centered practice to increase engagement

### 1. Billing and Coding

#### *Initiation in Primary Care*

Your patient must have **one or more** of these visit types **within 14 DAYS** of the initial diagnosis to meet the measure.

#### **For patients with all types of SUD:**

Type of Visit	Common Codes
<b>In-Person Office Visit with a SUD Dx</b> *Substance must match the member's initial Dx type	<b>E&amp;M Codes</b> 99211-99215, 99203-99205  <b>BH Services Integrated in Primary Care</b> 90971, 90972, 90832, 90837, 90840, 90847, 90849, 90853 H0001, H0002, H0031
<b>Telephone Visit with a SUD Dx</b> *Substance must match the member's initial Dx type	98966, 98967, 98968 99441, 99442, 99443
<b>E-Visit/ Virtual Visit with a SUD Dx</b> *Substance must match the member's initial Dx type	99421, 99423, 99444 G0071, G2012

#### **For patients with OUD (in addition to codes listed for all substance types):**

Type of Visit/Claim	Medication list
<b>OUD Medication Prescription</b>	<b>Naltrexone</b> (oral or injectable) <b>Buprenorphine</b> (sublingual tablet, injection, implant) <b>Buprenorphine/Naloxone</b> (sublingual tablet, buccal film, sublingual film)

#### **For patients with AUD (in addition to codes listed for all substance types):**

Type of Visit/Claim	Medication list
<b>AUD Medication Prescription</b>	<b>Disulfiram</b> (oral) <b>Naltrexone</b> (oral or injectable) <b>Acamprosate</b> (oral; delayed-release tablet)

### Engagement in Primary Care

A patient must have the right combination of visit types **within 34 DAYS of initiation** to be considered engaged in treatment. How a patient can become engaged depends on their type of treatment initiation:

Initiation Type	Meet criteria for 'Engaged' in Metric	
Patients who initiated treatment <b>WITH</b> medication	One Medication Event + One Engagement Visit	Two Engagement Visits
Patients who initiated treatment <b>WITHOUT</b> medication	One Medication Event	Two Engagement Visits

### Medication event in primary care setting

Type of Claim	Medication list
<b>ODU Medication Prescription</b>	<b>Naltrexone</b> (oral or injectable) <b>Buprenorphine</b> (sublingual tablet, injection, implant) <b>Buprenorphine/Naloxone</b> (sublingual tablet, buccal film, sublingual film)
<b>AUD Medication Prescription</b>	<b>Disulfiram</b> (oral) <b>Naltrexone</b> (oral or injectable) <b>Acamprosate</b> (oral; delayed-release tablet)

### Engagement visit in primary care:

Type of Visit	Common Codes
<b>In-Person Visit with an SUD Dx</b> *Substance must match the member's initial Dx type	<b>E&amp;M Codes</b> 99211-99215, 99203-99205  <b>BH Services Integrated in Primary Care</b> 90971, 90972, 90832, 90837, 90840, 90847, 90849, 90853 H0001, H0002, H0031
<b>Telephone Visit with an SUD Dx</b> *Substance must match the member's initial Dx type	98966, 98967, 98968 99441, 99442, 99443
<b>E-Visit/ Virtual Visit with an SUD Dx</b> *Substance must match the member's initial Dx type	99421, 99423, 99444 G0071, G2012

\*These codes count as a medication event AND an engagement event for members in the OUD cohort, so they meet the engagement metrics on their own.

## 2. BHC Services in Primary Care

Behavioral Health Consultants (BHC) can help create an environment in primary care that supports an open door for recovery. Ensure patients know that the BHC is a resource, the BHC should be introduced to patients who are diagnosed with a substance use disorder, receive medication for substance use, or receive a follow up from ED visit for substance use

### Key Services

- Preventative medicine counseling
- Psychotherapy
- Health and Behavior (for SBIRT)

### Key BHC Workflows

- Utilize BHC for SBIRT, this is the start of **initiation/identification**; BHC can screen during BHC and PCP appointments
- BHC can help facilitate referrals and coordinate care if outside referrals are the best course of treatment
- Introduce BHC to patients who are diagnosed with any use disorder

## 3. Collective Medical

**Real time knowledge of SUD inpatient and emergency department admissions allow us to coordinate in the moment to best meet our members' needs.**

- Set up alerts/notifications to know when your patients end up in the ED for SUD related issues so you can follow up quickly to provide support
- Consider utilizing Collective IET cohort (details in chart below)
- Utilize reports as an additional resource for scrubbing/reviewing records before visits
- Create watch lists of patients whom you've seen in the clinic for better monitoring
  - Include those who you're referring to behavioral health and those who are going to follow up with you in primary care
  - If patients go to ED for behavioral health related issues (substance use or not), reach out for quick follow up

## Collective Medical Cohort Criteria

The specifications below are suggested cohort criteria based on CareOregon's Collective onboarding support for health care providers. Some providers/organizations may choose to adjust criteria to best meet their organization's resources, needs, and existing workflows.

### SUD-IET—Any Encounter Event

- Triggering Event: *Any visit activity in ED, Inpatient, Observation settings*
- Physical Age above 18+
- Exclusions:
  - Discharge code does not equal 20 (to indicate Patient 'Expired')
  - F17 Nicotine related
- Diagnosis Phrase:  
REGEX-(?i)alcohol.\*|dependence.\*|withdrawal.\*|abuse.\*|drug use.\*|heroin.\*|opiate.\*|opioid.\*|adverse.\*|overdose.\*|tremens.\*|Intoxication.\*|poison.\*|hallucinogen.\*|Illicit.\*|detox.\*

### And/OR:

- Diagnosis Code:
  - Include the following ICD -10 codes with all subtypes if there is an asterisk:
  - F10\* Alcohol related
  - F11\* Opioid related
  - F13\* Sedative, hypnotic or anxiolytic related
  - F14\* Cocaine related
  - F15\* Other Stimulant related (this will capture methamphetamine use)
  - F16\* Hallucinogen related
  - F18\* Inhalant related
  - F19 Other Psychoactive substances
  - O9931\* / O9332\* Alcohol related, pregnancy Drug use complicating, childbirth, and the puerperium
  - T401\* Poisoning by Heroin
  - T402\* Poisoning by Opioid
  - T404\* Poisoning by synthetic narcotics
  - T409\* Poisoning by hallucinogens
  - T42 Poisoning by, adverse effect of and underdosing of antiepileptic, sedative- hypnotic and antiparkinsonism drugs
  - T51.91XA Toxic effect of unspecified alcohol, accidental (unintentional), initial encounter

#### 4. Person-Centered Best Practice

- Use a trauma informed, person-centered approach to educate and care for your patient
  - Use this opportunity to establish a supportive and trusting relationship with your patient with phrases such as “I’m so glad you are here”, “I care so much about your safety”
  - Language Matters! By using positive, person-centered language, you are more likely to keep people engaged in care. Feeling stigmatized can reduce the willingness of individuals with SUD to seek treatment.
- Prescribe naloxone for any person who has a substance use disorders; ensure your patient has naloxone in-hand
- Discuss Medication for Opioid Use Disorder (MOUD) with any person diagnosed with Opioid Use Disorder (OUD).
  - Prescribe MOUD when indicated.
  - MOUD is the gold-standard, best-practice for the treatment of OUD.
- Connect your patient with syringe access, wound care supplies, and community supports as needed
- Query the Prescription Drug Monitoring Program (PDMP) to identify all prescribers/prescriptions
  - Coordinate care with any outside providers
- Ask your patient to sign a Release of Information to access substance use treatment records

#### Links to Helpful Documents:

[Words Matter: Terms to Use and Avoid When Talking About](#)

[Addiction Recovery Oriented Language Guide](#)