

FIGURE 3–4c. CLASSIFYING ASTHMA SEVERITY IN YOUTHS ≥12 YEARS OF AGE AND ADULTS

■ **Classifying severity for patients who are not currently taking long-term control medications.**

Components of Severity		Classification of Asthma Severity (Youths ≥12 years of age and adults)			
		Intermittent	Persistent		
Impairment	Symptoms	≤2 days/week	>2 days/week but not daily	Daily	Throughout the day
	Nighttime awakenings	≤2x/month	3–4x/month	>1x/week but not nightly	Often 7x/week
	Short-acting beta ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week but not >1x/day	Daily	Several times per day
	Interference with normal activity	None	Minor limitation	Some limitation	Extremely limited
	Lung function	<ul style="list-style-type: none"> • Normal FEV₁ between exacerbations • FEV₁ >80% predicted • FEV₁/FVC normal 	<ul style="list-style-type: none"> • FEV₁ ≥80% predicted • FEV₁/FVC normal 	<ul style="list-style-type: none"> • FEV₁ >60% but <80% predicted • FEV₁/FVC reduced 5% 	<ul style="list-style-type: none"> • FEV₁ <60% predicted • FEV₁/FVC reduced >5%
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year (see note)	≥2/year (see note) →		
		← Consider severity and interval since last exacerbation. Frequency and severity may fluctuate over time for patients in any severity category. →			
		Relative annual risk of exacerbations may be related to FEV ₁			

- Level of severity is determined by assessment of both impairment and risk. Assess impairment domain by patient’s/caregiver’s recall of previous 2–4 weeks and spirometry. Assign severity to the most severe category in which any feature occurs.
- At present, there are inadequate data to correspond frequencies of exacerbations with different levels of asthma severity. In general, more frequent and intense exacerbations (e.g., requiring urgent, unscheduled care, hospitalization, or ICU admission) indicate greater underlying disease severity. For treatment purposes, patients who had ≥2 exacerbations requiring oral systemic corticosteroids in the past year may be considered the same as patients who have persistent asthma, even in the absence of impairment levels consistent with persistent asthma.

■ **Classifying severity in patients after asthma becomes well controlled, by lowest level of treatment required to maintain control.***

Lowest level of treatment required to maintain control (See figure 4–5 for treatment steps.)	Classification of Asthma Severity			
	Intermittent	Persistent		
	Step 1	Mild	Moderate	Severe
		Step 2	Step 3 or 4	Step 5 or 6

Key: EIB, exercise-induced bronchospasm; FEV₁, forced expiratory volume in 1 second; FVC, forced vital capacity; ICU, intensive care unit

*Notes:

- For population-based evaluations, clinical research, or characterization of a patient’s overall asthma severity after control is achieved. For clinical management, the focus is on monitoring the level of control (See figure 3–5c.), not the level of severity, once treatment is established.
- See figure 3–5c for definition of asthma control.